

National University Health System

Yong Loo Lin School of Medicine • National University Hospital • Faculty of Dentistry



Achieving Functional Cure in HBV

SHC-HK ASLD Webinar series


**Prof Seng Gee Lim
Director of Hepatology,
Dept of Gastroenterology and Hepatology
National University Health System
Singapore**




Disclosures

- Advisory Board
 - Gilead
 - Roche
 - Abbvie
 - Abbott
 - Springbank
 - Kaleido Bioscience
- Speaker's Bureau
 - MSD
 - Gilead
 - Abbvie
 - Abbott

Endpoints versus Goals



Endpoints in HBV Treatment



What is HBV Functional Cure?


Term	Definition	Achievability
Functional Cure	• Sustained HBsAg loss/seroconversion 24 weeks off therapy • Undetectable HBV DNA	• Can be achieved with existing therapy • Ideally ~30% in phase III studies
Absolute or complete Cure	• Absence of cccDNA • Absence of integrated HBV	• Unlikely to be achieved in foreseeable future

HBsAg loss is associated with improved survival & complications: meta analysis


Endpoint	n ^{srcr}	P-Y ^{srcr}	n ^{prstnt}	P-Y ^{prstnt}	RR and 95% CI	Reduction	RR	Lower limit	Upper limit	P-value
Liver Decompensation	4382.7	21426	105126.3			72%	0.28	0.13	0.59	.001
HCC	273264.3	154519	1211810.9			70%	0.30	0.20	0.44	<.001
LTx/death	7663.9	9131	69053.6			78%	0.22	0.13	0.39	<.001
First Clinical Event	273095.2	155030	1212985.9			69%	0.31	0.23	0.43	<.001

Variable	Subgroup	K	N	P-Y		RR	Lower limit	Upper limit	P-value
Study Design									
	Retrospective	18	182465	1416301.5		0.27	0.17	0.44	<.001
	Prospective	10	5851	69779.7		0.33	0.19	0.55	<.001
Cohort type									
	Untreated	12	155534	1312287.0		0.30	0.21	0.44	<.001
	Treated	13	31086	151408.5		0.21	0.08	0.54	.001
	Both	3	1696	22385.6		0.33	0.15	0.75	.008
Race									
	Caucasian	9	2698	18861.3		0.24	0.12	0.48	<.001
	Asian	18	185180	1461423.9		0.28	0.18	0.46	<.001
	Other	1	438	10849.0		0.67	0.25	1.85	.444
Co-infected subpopulation									
	Yes	3	148485	1217211.2		0.31	0.16	0.61	.001
	No	25	39831	268869.9		0.31	0.21	0.45	<.001
HBsAg Status at Baseline									
	(+)	7	1717	10496.7		0.22	0.08	0.64	.005
	(-)	6	3787	45751.3		0.22	0.10	0.48	<.001
	Both	15	182812	1429833.2		0.31	0.18	0.51	<.001
									
					0.0Overall*0.51.01.5				
					RR				


Factors leading to HBsAg loss




Feature	Subgroup	K	N	ASR (95% CI)
Sex	Male	16	11142	1.13 [0.78; 1.55]
	Female	14	6080	0.87 [0.49; 1.36]
Baseline HBsAg	HBsAg+	9	3922	0.40 [0.25; 0.59]
	HBsAg-	9	15205	1.33 [0.76; 2.05]
Baseline cirrhosis	No	12	13398	1.04 [0.63; 1.55]
	Yes	8	970	0.84 [0.31; 1.58]




Feature	Subgroup	K	N	ASR (95% CI)
Overall		34	42588	1.02 [0.79; 1.27]
Study region	North America	3	4984	0.50 [0.15; 1.04]
	Europe	5	2364	1.08 [0.64; 1.63]
	Asia Pacific	23	32866	1.09 [0.81; 1.42]
	Middle East	2	1826	1.05 [0.82; 1.31]
	South America	1	548	0.70 [0.50; 0.94]
Practice setting	Community	8	9435	1.34 [0.92; 1.83]
	University	16	9521	0.97 [0.78; 1.18]
	Mixed	6	12101	0.74 [0.16; 1.71]
Scale of study	Population-based	4	5779	1.68 [0.86; 2.78]
	Healthcare service-based	30	36809	0.93 [0.72; 1.17]



Variable	HBsAg seroclearance	K	N	Mean (95% CI)
qHBsAg (log ₁₀ IU/mL)	Yes	5	1173	2.74 [1.87; 3.60]
	No	5	6246	3.90 [3.73; 4.06]
ALT (U/L)	Yes	16	1663	63.75 [27.09; 149.99]
	No	16	14974	58.18 [19.08; 177.46]
HBV DNA (log ₁₀ IU/mL)	Yes	10	1431	6.61 [5.94; 7.27]
	No	10	16590	7.71 [7.41; 8.02]
Platelet count (10 ⁹ /L)	Yes	10	822	195.14 [182.82; 208.29]
	No	10	9536	195.71 [182.02; 210.43]



What approaches can we use with current therapies?




Which is the best strategy?


HBsAg loss with current therapy: meta analysis

43 high quality RCTs with low risk of bias


Combination(IFN+NA) vs NA



IFN vs NA



Combination(IFN+NA) vs IFN



Combination

NA

6.55%

0.58%

IFN

NA

3.64%

0.27%

Combination

IFN


4.8%

3.6%

- IFN or combination is superior to NA but no difference to each other
- Current therapies achieve HBsAg loss < 10%

Patterns of qHBsAg decline during 5y TDF therapy


Modelling predicts 36-39y of NUCs to achieve HBsAg loss



HBsAg loss - Genotype A	N = 14	14	14	13	13	13	13	12	12	11	10	11	10	10	9	8	8	7	7
HBsAg loss - Genotype B	N = 1	1	1	1	1		1	1	1	1	1	1	1	1	1	1	1		
HBsAg loss - Genotype D	N = 7	7	7	7	7	6	6	6	6	6	4	4	3	2	2	3	2	2	
No HBsAg loss - Genotype A	N = 45	44	43	43	42	40	40	40	40	39	39	35	38	37	36	37	35	32	33
No HBsAg loss - Genotype B	N = 34	33	33	31	30	26	24	24	22	23	22	22	22	22	22	21	22	21	21
No HBsAg loss - Genotype C	N = 68	68	66	65	65	60	59	57	57	58	58	57	56	53	53	52	51	47	46
No HBsAg loss - Genotype D	N = 77	76	77	73	73	72	68	67	66	65	62	62	59	58	58	57	55	58	53

Stopping NA therapy

What can we learn from stopping NA therapy?



Increased T cell activation and reduced exhausted T cell phenotype in those who stop therapy

Stopping Long-Term Nuc Therapy

	Stay off treatment	HBsAg loss	Predictors of HBsAg loss	PPV	NPV
Sonneveld (48 wk F/U)	53%	4%	BL HBsAg <10	27% (7/26)	98% (428/43)
van Bömmel (48 wk F/U)	86%	10%	BL HBsAg<1000	27% (7/25)	98% (53/54)
Hall (48 wk F/U)	89%	5%	BL HBsAg <10	63% (5/8)	100% (95/95)


- NUC discontinuation may be safe in noncirrhotic patients
- 30-60% will remain off-treatment and 5-10% will lose HBsAg
- Best predictor is low HBsAg level at discontinuation

Likelihood of HBsAg loss after stopping NA therapy based on qHBsAg levels: Systematic Review

EOT qHBsAg level	HBsAg loss
<100 IU/mL	21.1%-58.8%
100-200 IU/mL	25.5%
100-500 IU/mL	4.4%
>500 IU/mL	2.3%
>1,000 IU/mL	0.0%


**After HBsAg loss, can HBV
return?**

HBV reactivation in HBsAg negative, anti-HBc Ab positive patients with Rituximab Therapy




- Blocking B cell function leads to HBV reactivation in ~15% of HBsAg negative patients


De novo HBV post Liver Transplant



Liver Transplant
Donor



HBsAg neg
Anti-HBc pos



Liver Transplant
Recipient


**Up to 72% De Novo HBV
In anti-HBc neg recipients**

Summary

- Immune suppression leads to reactivation of HBV in those who are HBsAg neg, anti-HBc positive
- In the most extreme case of liver transplant this is as high as 72%
- Indicates that cccDNA has not been eliminated
- More likely cccDNA has been silenced
- Measurement of cccDNA in liver tissue as an endpoint of therapy may be problematic
 - Sensitivity not high
 - Assays not standardized

**New Therapies are needed
to achieve functional cure**


Novel therapeutic targets for HBV




CpAM: core protein allosteric modulators; HBx: hepatitis B X protein; IFN: interferon; IL: interleukin; KC: Kupffer cells; mAb: monoclonal antibody; NA: nucleos(t)ide analogue; NK (T): natural killer (T cell); DC: plasmacytoid dendritic cell; PD-1: programmed cell death-1; TCR: T cell receptor

Adapted from: Leviero M, et al. Curr Opin Virol 2016;18:135–43;

Three Strategies



Overlapping pathways in HBV lifecycle



Two main controllers of viral transcription

- Core promoter & enhancers control pregenomic RNA
- **"S" promoter controls "S" transcripts**
- S Ag production is affected by amount of cccDNA and its activity, as well as by epigenetic factors controlling "S" promoter (eg methylation)

Functional Cure

Translational Pathway

Replicative Pathway

How do we know which agents are effective?

HBsAg


- HBsAg loss ideal but difficult to achieve in Phase I-II RCT
- qHBsAg < 100
- Reduction in qHBsAg (>1 log, >2 log?), the bigger the better

Other markers

- HBeAg loss/seroconversion
- HBV DNA reduction
- HBV RNA reduction
- HBcrAg reduction
- Measurement of cccDNA

Biomarkers for HBsAg loss in CHB treated with pegIFN+NA

▲ mean HBcrAg log IU/mL ■ mean qHbsAg log IU/mL ● mean HBV RNA log IU/mL



Baseline AUROC for HBsAg loss


- qHBsAg (0.916)
- HBcrAg (0.649)
- HBV RNA (0.542)

On treatment qHBsAg week 8 was best predictor


- Threshold 69 IU/ml
- AUROC=0.963

qHBsAg was superior to HBV RNA and HBcrAg in predicting HBsAg loss

Definitions of response




Pathways that inhibit/reduce intracellular HBV & replication




CpAM: core protein allosteric modulators; HBx: hepatitis B X protein; IFN: interferon; IL: interleukin; KC: Kupffer cells; mAb: monoclonal antibody; NA: nucleos(t)ide analogue; NK (T): natural killer (T cell); DC: plasmacytoid dendritic cell; PD-1: programmed cell death-1; TCR: T cell receptor

Capsid Assembly Modulators (CAMs)



Antiviral Activity, Safety, and PK of ABI-H0731 in CHB (Study101B) without NA




Study Design:



- Once Daily Dosing
- HBeAg (+), (-): 7:5
- Follow up for 28 days



Efficacy




Safety





Adverse Event	100mg (N=10)	200mg (N=10)	300mg (N=10)	400mg (N=2)	Placebo (N=6)
Grade 1	9 (90%)	4 (40%)	6 (60%)	2 (100%)	4
Grade 2	0	0	0	0	0
Grade 3	0	0	0	1 (Rash)	0
ALT Flare	No ALT flare was observed during or post treatment				

Treatment with 24 weeks CAM +NA

- Phase II JADE study of JNJ-6379 (CAM-N, class II)
 - 2 doses (75mg and 150mg) plus NUC
 - Monotherapy arms d/c due to breakthrough
 - Read out at 24 weeks
 - Primary endpoint is change in HBsAg level



^aJNJ-6379 treatment could be extended to 48 weeks if response criteria were met.
 Red boxes indicate this presentation focused on placebo and JNJ-6379 arms only




Duration of Treatment (weeks)

Treatment with 72 weeks CAM +NA


- Phase 2 open label extension study of Vebicorvir (ABI-H0713)
- Class II CAM






1. HBeAg negative CHB¹



2. HBeAg positive CHB²



- Efficacy: HBsAg decline at end of 72 weeks VBR



¹ Fung S, et al. EASL dILC2020. #AS070

² Yuen M-F, et al. EASL dILC2020. #LBP30


Efficacy of CAMs

Compound	Pharma	class	CHB	Duration & therapy	HBV DNA	HBV RNA	HBsAg
JNJ-56136379	Janssen	II	Not treated	24w +NA	↓4-5.89 log	↓3.13 log	↓0.4 log in HBeAg(+)
JNJ-56136379		II	Treated	24w +NA	NA	↓1.07 log	No change
ABI-H0731	Assembly	II	Not treated	75w +NA	50% LLOQ	NA	40%≥0.5 log↓
ABI-H0731		II	Treated	77w +NA	NA	NA	2%≥0.5 log↓
AB-506	Arbutus	II	Not treated	28d	↓2.8 log	↓2.4 log	No change


Janssen, LBP-012, EASL 2020
Sims, HepDART 2019
Yuen, EASL 2020

Triple therapy: NA+CAM+RNAi

HBsAg



HBV DNA





- HBeAg+ n=4 / HBeAg- n=8, NA-naïve n=5 / experienced n= 7, All 12 Asian
- Three 200 mg JNJ-3989 subcutaneous doses on Days 1, 29 and 57
- Oral JNJ-6379 250 mg once daily for 12 weeks (until Day 85)
- Started or already on ETV or TDF treatment on Day 1 to beyond the end of JNJ-6379 dosing
- Response rates similar between HBeAg+ and HBeAg-

Summary


- CAMs are a new class of antiviral agents that disrupt capsid formation
- Early phase studies show reduction in HBV DNA and pgRNA but little change in qHBsAg levels
- Extended duration studies show that some patients may have reduction in qHBsAg
- Unlikely it can be used alone, more likely in combination

Entry inhibitors: Myrcludex







Myrcludex B is a synthetic lipopolypeptide consisting of a myristoylated pre-S1 domain of the large HBsAg with binding to the HBV entry receptor


MYR203 Study for HBV-HDV co-infection



- 60 patients with chronic HBV/HDV co-infection were randomized into 4 treatment arms in a ratio of 1:1:1:1 - 15 patients per arm
- MyrB was self administered by patients once daily s.c.




Definition of HBsAg Response ($\geq 1\log_{10}$ decline or undetectable)



Summary

- Myrcludex B monotherapy (2mg sc/qd):
 - continuous linear HDV RNA decline and ALT reduction over 48 weeks
 - relapse in most patients (73.3%) after the end of therapy
 - However **no significant reduction in qHBsAg**
- Myrcludex B/PEG-IFN α combination therapy:
 - HDV RNA undetectable in 53.3% (2mg/PEG-IFN α) and 26.7% (5mg/PEG-IFN α) at week 72, and 0% (PEG-IFN α)
 - HBsAg response in 40% (2mg/PEG-IFN α ; incl. **26.7% HBsAg loss**) and 13.3% (5mg/PEG-IFN α) at week 72


Pathway to reduce viral antigen production



CpAM: core protein allosteric modulators; HBx: hepatitis B X protein; IFN: interferon; IL: interleukin; KC: Kupffer cells; mAb: monoclonal antibody; NA: nucleos(t)ide analogue; NK (T): natural killer (T cell); DC: plasmacytoid dendritic cell; PD-1: programmed cell death-1; TCR: T cell receptor


Adapted from: Leviero M, et al. Curr Opin Virol 2016;18:135–43;

RNA interference: ASO vs siRNA



Anti-Sense Oligo/LNAs

- RO7062931
- GSK3228836
- GSK3389404
- ISIS 505358




siRNAs

- ARC-520
- ARC-521
- ALN-HBV
- ARB-1467
- ARB-1740
- AB-729
- ARO-B/JNJ-3989
- GalXC-HBVS/DCR-HBVS
- ALN HBV02/VIR-2218

Mechanism of RNA interference: block production of viral proteins in a sustained manner esp HBsAg, hence reduce viral antigen burden, and HOPEFULLY allow recovery of immune cell exhaustion (not yet been demonstrated)

RO7062931: a liver-targeted, single-stranded oligonucleotide (SSO) with locked nucleic acid (LNA)

- RO7062931 is an SSO LNA targeting a highly conserved sequence in the HBV genome
- N-acetylgalactosamine(GalNAc) conjugation achieves liver-targeting through interaction



- RO7062931 treatment resulted in statistically significant* time-dependent HBsAg reductions compared to placebo for the 1.5 mg/kg and 3.0 mg/kg dose groups
- Maximum HBsAg decline typically occurred ~2wks post-treatment, with rebound to baseline levels by ~12wks post-treatment
- HBV RNA levels showed similar, although less profound and more variable, kinetics (data not shown)
- No significant changes were observed for HBcrAg levels (data not shown)

*90% Confidence Intervals for adjusted treatment differences from the statistical analysis (data not shown) did not include 0.


HBcrAg, hepatitis B core-related antigen

8


Short term therapy leads to reduction in qHBsAg<0.5 log followed by rebound

AB-729 RNAi Therapeutic


- Single trigger RNAi agent targeting all HBV transcripts
- Inhibits HBV replication and lowers all HBV antigens
- Pan-genotypic activity across HBV genotypes
- Demonstrated complementarity with capsid inhibitors
- GalNAc-conjugate actively targets the liver
- Active against cccDNA derived and integrated HBsAg transcripts




AB-729 60 mg (N=6)



AB-729 90 mg (N=6)




AB-729 180 mg (N=4)



siRNA JNJ-3969 (ARO-HBV)

- 3 X Q4W doses of JNJ-3989 + NUC in CHB patients
- Extended follow-up to 1 year post-siRNA to assess long-term safety of siRNAs and also sustained responses in HBsAg, HBV RNA, HBeAg and HBcrAg

Cohort	N	Dose
2b	8	3 x 100 mg Q4W
3b	8	3 x 200 mg Q4W
4b	8	3 x 300 mg Q4W
5b	8	3 x 400 mg Q4W
8 eAg pos, Nuc-naïve	4	3 x 300 mg Q4W
9 eAg pos, Nuc-exp	4	3 x 400 mg Q4W






janssen Infectious Diseases & Vaccines
PHARMACEUTICAL COMPANIES OF Johnson & Johnson

Sustained off therapy response after almost 1y

- Efficacy: HBsAg Response (n=40)**
- 85% had ≥ 1 log reduction during treatment.
- 39% had ≥ 1 log reduction at 48 weeks post-dose; i.e. were sustained HBsAg responders
- Off-Treatment Sustained HBsAg Responder**
- ≥ 1 Log₁₀ IU/mL reduction in HBsAg from at Day 392

ASO ISIS 505358 (GSK)

- Naked (non-Gal-NAC) conjugated ASO
 - 300mg Loading and frequent dosing over 28 days
 - NUC-naïve (12) and suppressed (4); HBeAg pos and neg



- 4 patients had levels falling below the limit of quantification (qHBsAg<0.05 IU/ml).
- **Prolonged HBsAg loss was observed in 2 patients:**
 - one-NA treated patient (from Day 36 to Day 113)
 - one NA-naïve patient (from Day 23 to Day 126).


Summary

- RNA interference is directed to reduce production of viral proteins, esp HBsAg
- Needs to be given by injection
- Studies of extended followup after limited treatment show sustained suppression of HBsAg for over one year
- Proof of sustained HBsAg loss demonstrated
- There have been associated ALT flares with reduction in qHBsAg
- Most likely it will be best used in combination therapy


Nucleic Acid Polymers (NAPs)

S-antigen traffic-inhibiting oligonucleotide polymers (STOPs)


Company	Investigational New Drug	Clinical Trial
Replicor Inc.	REP-2139	Phase 2
Replicor Inc.	REP-2165	Phase 2
Replicor Inc.	REP-2055	Phase 1



Bazinet M, EASL-2017, #LBP-507



Efficacy results with REP 9AC monotherapy



REP 2055 was given by IV infusion 400mg weekly for 40w

Number of patients	Number with sustained off therapy responses	No with viral relapse req rescue or re-treatment
8 HBeAg negative CHB	2 (25%)	5*

* One patient lost to followup

Nucleic Acid Polymers (REP-2139, 2165): Study 401


REP 401 is a randomized, controlled trial assessing safety and efficacy of REP+TDF+PEGIFN in HBeAg (-) CHB.

IND	NAP Sequence Design	Stable
REP-2139	ACACACACACACAC ACACACACACACAC ACACACACACAC	93%
REP-2165	ACACACACACAC* ACACACACACAC ACACACACACAC	89%

Study Design (250mg, IV, QW)					
Week	0	24	48	72	96
eAg (-) N=20	TDF	TDF PEG-IFN REP	TDF PEG-IFN REP		
eAg (-) N=20	TDF	TDF PEG-IFN ★ REP	TDF PEG-IFN REP	TDF PEG-IFN REP	

BL Factor	Study Arm	Ctrl Arm
Age (m/m)	36.9/36.0	38.6/39.5
Sex (M/F)	17/3	16/4
GT (A/D)	1/19	2/18
F (1/2/3/4)	12/4/0/4	10/7/2/1
HBsAg/Ab (IU/mL)	9300/0.1	8700/0.1
HBV DNA (IU/mL)	3.6x10 ⁷	4.8x10 ⁶

*Replacement of 2'Ome with 2'OH at 11, 21 and 31



Final REP 401 outcome summary

Completed treatment and ≥ 24 weeks of follow-up		35 (32 completed 48 weeks of follow-up)
Clinical response	Normal ALT	91%
	Normal median hepatic stiffness	57%
HBsAg response	< 1000 IU/mL	74%
	< 1 IU/ml	51%
	$\leq$ LLOQ (0.05 IU/mL)	43%
	Seroconversion	54%
HBV DNA response	≤ 2000 IU/mL	83%
	< 100 IU/mL	63%
	Target not detected (TND)	49%
Virologic response	Functional control (HBV DNA ≤ 2000 IU/mL, normal ALT)	43%
	Functional cure (HBsAg $< \text{LLOQ}$, HBV DNA TND, normal ALT)	40%
	Clinical benefit, no therapy required (Low risk of progression, reduced risk of HCC)	83%


Summary

- This novel treatment shows promise even as monotherapy
- Impact of TDF+pegIFN has limited impact on HBsAg reduction until addition of NAP
- Combination therapy with NA and pegIFN has demonstrated functional cure in 40%

Immune Modulation


Therapeutic Vaccines

AIC 649	AiCuris, Germany
INO-1800	Inovio, USA
HB-110	Ichor Med Systems with Janssen, USA
TG1050	Transgene, France
HepTcell	Altimmune, USA
HB-110	Ichor Medical Genexine, USA
VTP-300	Vaccitech, USA




CpAM: core protein allosteric modulators; HBx: hepatitis B X protein; IFN: interferon; IL: interleukin; KC: Kupffer cells; mAb: monoclonal antibody; NA: nucleos(t)ide analogue; NK (T): natural killer (T) cell; DC: plasmacytoid dendritic cell; PD-1: programmed cell death-1; TCR: T cell receptor

Overview of immune modulation therapies for HBV




Toll-like Receptor Agonist Mechanism of Action

- TLRs are PRRs that recognise pathogen-associated molecular patterns
- Crucial for early host response and linking innate with adaptive responses



IFN, interferon; IL, interleukin; MAIT, mucosal-associated invariant T cells; mDC, myeloid dendritic cells; NK, natural killer cells; pDCs, plasmacytoid dendritic cells; PRR, pattern recognition receptor; TNF, tumor necrosis factor; Tregs, regulatory T cells. 1. Schurich A, et al. PLoS Pathog 2013;9(3):e1003208; 2. Du H, et al. Vaccine 2013;28;31:4209-15; 3. Jo J, et al PLoS Pathog 2014;10:e1004210; 4. Isorce N, et al. Antiviral Res 2016;130:36-45.



Activation of Innate Immunity in HBV



Efficacy and Safety of Oral TLR8 Agonist Selgantolimod (GS-9688, SLGN) in Virally Suppressed Adult Patients with CHB: Phase 2 study

- Primary endpoints:

- Safety and tolerability of selgantolimod at Week 24
- Proportion of patients with $\geq 1.0\text{-log}_{10}$ IU/mL HBsAg decline from baseline at Week 24




In SLGN-treated patients, HBsAg was sustained or continued to decline during the 24 weeks of post-treatment follow up HBsAg decline $\geq 0.1\text{-log}_{10}$ IU/mL was observed only in SLGN-treated patients

- 2/39(5%) patients had HBsAg loss
- However the reduction in qHBsAg overall was <1 log
- One patient with HBsAg loss had only ~ 0.25 log reduction


Adaptive immunity: restoration of exhausted T cells: PD-1 inhibition

Nivolumab: Monoclonal antibody against PD-1 Approved for solid organ tumors and lymphomas

GS-4774: Heat-inactivated yeast-based therapeutic T-cell vaccine developed for CHB



1/10 patients had HBsAg seroclearance




0/12 patients had HBsAg seroclearance

Gane, EASL 2017

Therapeutic Vaccines: can they work?

Principle: Develop new vaccines against HBV by targeting HBsAg or HBcAg using novel adjuvants or vectors




Problem: overcoming immune tolerance to HBV due to huge HBsAg antigen load

Meta-analysis of RCTs: Therapeutic Vaccine vs No Therapy/Placebo

HBeAg seroconversion

[End of followup (off therapy)]




Total 16 RCTs

- Therapeutic Vaccine=1090 pt
- PLB/no therapy=591 pt

HBsAg loss

[End of followup (off therapy)]



Therapeutic Vaccine vs no therapy/placebo = No efficacy for:


- HBeAg SC
- HBsAg loss

Meta-analysis: Therapeutic Vaccine + SOC* vs SOC*

HBeAg seroconversion


[End of followup (off therapy)]

*SOC = Standard of Care



HBsAg loss

[End of followup (off therapy)]



Therapeutic Vaccine + SOC vs SOC

= No efficacy for:

- HBeAg SC
- HBsAg loss

What type of studies have led to significant functional cure?

Interferon based studies

- PegIFN mono or Nuc/PegIFN therapy 8 - 10% (End of FU)
- Myrcludex B 2mg/pegIFN 48w in HBV/HDV co-infection 26.7% (End of FU)
- TDF + pegIFN + NAP Triple Therapy 40% (End of FU)

Stopping Nucs studies

- 19-46% HBsAg loss from different studies
- Rates for HBsAg loss rely on low qHBsAg at EoT

Hadziyannis SJ, et al. *Gastroenterology*. 2012;143:629-636
Papatheodoridis, *Antiviral Ther* 2018;23(8):677-685
Chen, *Clin Microbiol Infect*. 2018 Sep;24(9):997-1003
Yao, *Sci Rep*. 2017 May 12;7(1):1839

More than one path to functional cure


Potent combination therapy

- Combine agents from different classes
- Most of these combinations rely on immune modulators (48w therapy)
 - JNJ triple combination
 - Mycludex, TDF + pegIFN (26.7% HBsAg loss)
 - NAPs, TDF +pegIFN (40% HBsAg loss)

HBsAg reduction followed by immune therapy?

- Stopping NA successfully leads to functional cure in qHBsAg<100 IU/ml associated with HBV flares and improved T cell function, but HBsAg loss takes as long as 3-5years
- Potential therapeutic strategy

HBV Treatment Strategies



Conclusions

- Functional cure is an achievable endpoint associated with improved clinical outcomes
- Functional cure is most likely associated with cccDNA silencing than eradication
- Agents that have little effect on qHBsAg in early studies cannot be completely discounted (eg Myrcludex), and may still have efficacy in combination
- Combination therapy is the most promising but sequential therapy may be another therapeutic strategy.
 - This may be useful when agents can reduce qHBsAg < 100 IU/ml, and given immunomodulators
- Quite likely there are multiple pathways to achieve functional cure, but therapies currently that have been successful in achieving functional cure in >20% have relied on pegIFN

Acknowledgements

Special thanks to the following for allowing their slides to be shared

- Gilead
- Janssen
- Roche
- Assembly
- Arbutus
- Replicor
- Ed Gane



Sharp focused updates in Hepatology

UPDATES

3 ADDITIONAL LIVE WEBINARS

Topic: Speaker: Date And Time:	New Biomarkers For Chronic Hepatitis B: Ready For Clinical Use? PROF. M.F. YUEN Monday, 12 October 2020 8.00pm – 9.00pm HKT/SGT	
Topic: Speaker: Date And Time:	Unmet Needs In CHB Guideline PROF. MARC GHANY Monday, 19 October 2020 8.00pm – 9.00pm HKT/SGT	
Topic: Speaker: Date And Time:	Should We Stop Long Term Nucleoside Analogue Therapy? PROF. TEERHA PIRATVISUTH Monday, 26 October 2020 8.00pm – 9.00pm HKT/SGT 7.00pm - 8.00pm Bangkok Time	

The logo for the Singapore Hepatology Conference is a red and white arrow pointing to the right. Inside the arrow, the words "Singapore", "Hepatology", and "Conference" are stacked vertically in a blue, sans-serif font. The word "Singapore" is at the top, "Hepatology" is in the middle, and "Conference" is at the bottom.

Singapore Hepatology Conference

Sharp focused updates in Hepatology

PLEASE REMEMBER TO FILL IN THE SURVEY FORM!